

**10% OF  
A BROCHURE  
MAY NOT BE  
ENOUGH**

**IF YOU HAVE IRON  
DEFICIENCY ANEMIA  
(IDA) DUE TO RECENT  
PREGNANCY OR  
HEAVY PERIODS...**

**10% OR LESS  
OF ORAL IRON  
MAY NOT BE  
ENOUGH EITHER**



Actor Portrayal.

Injectafer® (ferric carboxymaltose injection) is available by prescription only. Ask your doctor or healthcare provider if Injectafer is right for you.

## **What is Injectafer?**

Injectafer is a prescription iron replacement medicine administered only by or under the supervision of your healthcare provider. Injectafer is injected into your vein to treat iron deficiency anemia in adults and pediatric patients 1 year of age and older. Injectafer should be used only if you have not responded well to treatment with oral iron, or if you are intolerant to oral iron treatment. It is also used to treat iron deficiency anemia in adults with chronic kidney disease who are not receiving dialysis.

## **SELECTED SAFETY INFORMATION**

### **Who should not receive Injectafer?**

**You should not receive Injectafer if you** are allergic to ferric carboxymaltose or any of the other ingredients in Injectafer. The active ingredient in Injectafer is ferric carboxymaltose, the inactive ingredients are: water for injection, sodium hydroxide and/or hydrochloric acid.

**[Click here](#) to see the Full Prescribing Information and the Important Safety Information on pages 13-15.**



# GET TO KNOW INJECTAFER, AN IV IRON TREATMENT FOR IRON DEFICIENCY ANEMIA (IDA)



## SELECTED SAFETY INFORMATION ABOUT INJECTAFER

### What is Injectafer?

Injectafer is a prescription iron replacement medicine administered only by or under the supervision of your healthcare provider. Injectafer is injected into your vein to treat iron deficiency anemia in adults and pediatric patients 1 year of age and older. Injectafer should be used only if you have not responded well to treatment with oral iron, or if you are intolerant to oral iron treatment. It is also used to treat iron deficiency anemia in adults with chronic kidney disease who are not receiving dialysis.

## Understanding IDA

Causes, symptoms, and diagnosis of this potentially serious, but treatable condition many women of childbearing age face.

## Treating IDA:

### Oral Iron and Injectafer

Challenges with oral iron supplements and how another treatment option—an intravenous (IV) iron—might help.

## Treatment With Injectafer

Where, when, and how Injectafer is administered, along with helpful tips on how to plan for infusions.

## Savings & Support

Find out if you're eligible and learn how to enroll.

**STAY INFORMED ABOUT IDA AND INJECTAFER:  
SIGN UP FOR EMAILS AND TEXT MESSAGES  
AT [INJECTAFER.COM/SIGNUP](https://www.injectafer.com/signup)**

**[Click here](#) to see the Full Prescribing Information and the Important Safety Information on pages 13-15.**

# WHAT IS IDA?

- **Iron deficiency anemia (IDA)** is a condition that interferes with the formation and function of red blood cells
- **Red blood cells contain hemoglobin**, which is a protein that helps carry oxygen from the lungs to cells in the body
- When you don't have enough healthy red blood cells to carry enough oxygen to cells throughout the body, that's **anemia**

## ONE IN FIVE WOMEN OF CHILDBEARING AGE HAS IDA

Women may be at risk for IDA due to the body's **increased demand for iron, or iron depletion through blood loss.**

### IDA MAY RESULT FROM



Pregnancy, childbirth, and/or postpartum



Long or abnormal menstrual periods

When iron is depleted or not fully replenished, **low iron levels** may occur, impacting your overall health. If iron deficiency progresses, it may cause **IDA, as iron helps produce hemoglobin.**

### SELECTED SAFETY INFORMATION ABOUT INJECTAFER

#### Who should not receive Injectafer?

**You should not receive Injectafer if you** are allergic to ferric carboxymaltose or any of the other ingredients in Injectafer. The active ingredient in Injectafer is ferric carboxymaltose, the inactive ingredients are: water for injection, sodium hydroxide and/or hydrochloric acid.

### WHAT ARE SOME OF THE SYMPTOMS OF IDA?\*



Fatigue



Weakness



Dizziness



Pica  
(craving nonfood items such as ice or dirt)



Shortness of breath

\*Some people with IDA may not experience symptoms. Injectafer has not been studied or approved for treating symptoms of IDA.

### HOW IDA IS DIAGNOSED AND MONITORED

Certain lab markers in your blood are used to determine whether your iron levels are within the range your doctor feels is normal for you. These values can be used to diagnose and monitor IDA while being treated.

### COMMON IDA LAB MARKERS

#### NORMAL VALUES IN ADULT WOMEN†



<b>Hemoglobin (Hb)</b>	12.0-15.5 g/dL
<b>Ferritin</b>	20-200 µg/L
<b>Transferrin saturation (TSAT)</b>	20% to 50%

If your lab values are below the normal range, talk with your doctor about the best option for restoring your iron levels.

The information provided by Daiichi Sankyo, Inc., is informational only and not meant to provide medical advice.

An additional test for total iron binding capacity (TIBC) values may be required to diagnose IDA. Normal values for TIBC range from 240 to 450 µg/dL in healthy patients. Please note that TIBC values were not measured in Injectafer clinical trials.

†Normal values can vary for many reasons, including conditions you may have or where your lab work was done. That's why it's important to discuss your lab results with your doctor to find out what they mean for you.

**Click here to see the Full Prescribing Information and the Important Safety Information on pages 13-15.**



# HOW IS IDA TREATED?

## TWO OPTIONS TO TREAT IDA ARE:



Oral iron supplements



IV iron

## WHAT YOU SHOULD KNOW ABOUT ORAL IRON

- Oral iron supplements can cause hard-to-tolerate **side effects**
- **Poor absorption:** the digestive tract is only able to absorb a small portion of the iron in an oral iron supplement, so your body may not get the full dose of iron needed from pills. Even in healthy patients, **less than 10% of oral iron is absorbed**
- If you're being treated for IDA with oral iron supplements, talk to your doctor about checking your iron levels **within 14 and 30 days** of starting oral iron



Actor Portrayal.

**IF ORAL IRON  
ISN'T WORKING FOR  
YOUR IDA, IV IRON  
MAY BE AN OPTION**

## WHAT YOU SHOULD KNOW ABOUT IV IRON

- With IV treatment (known as infusions), **iron is delivered into the bloodstream** through a vein
- Since IV iron isn't processed by the digestive system, **100% is delivered into the bloodstream**

# INJECTAFER PROVIDES MORE IRON IN LESS TIME VS ORAL IRON

- For patients with iron deficiency anemia who **can't tolerate oral iron** or who **don't respond well to it**
- Injectafer is the only **IV iron** that provides up to **1500 mg of iron in one course of treatment\***

\*For patients weighing less than 50 kg (110 lb), the recommended dosage is Injectafer 15 mg/kg body weight intravenously in two doses separated by at least 7 days per course.



**1st administration**  
750 mg of iron

**2nd administration**  
750 mg of iron

**Injectafer is well studied and widely used, proven** in clinical studies with **more than 8800 people** and used to **treat more than 1.7 million** people in the United States alone.

**TALK WITH YOUR DOCTOR TO  
LEARN MORE ABOUT INJECTAFER**

## SELECTED SAFETY INFORMATION ABOUT INJECTAFER

### What should I tell my doctor or healthcare provider before receiving Injectafer?

Before you receive Injectafer, tell your healthcare provider about all of your medical conditions, including if you:

- Have had an allergic reaction to iron given intravenously (into your vein), including Injectafer, or to other non-oral iron treatments
- If you have, or have previously experienced, iron overload, or if your body has difficulty using iron appropriately
- Have high blood pressure

**Click here to see the Full Prescribing Information and the Important Safety Information on pages 13-15.**



# INJECTAFER INFUSIONS AT A GLANCE

Your OB/GYN or family doctor has prescribed Injectafer for you because it may be an effective way to replace the iron you need.

## WHERE INJECTAFER IS GIVEN

- **Injectafer is usually given at an infusion center** and administered by a healthcare professional
- Infusion centers are medical facilities equipped and staffed for **administering infusions**

**FIND AN INFUSION CENTER THAT PROVIDES INJECTAFER BY USING THE INJECTAFER INFUSION CENTER LOCATOR AT [INJECTAFER.COM/IRON-INFUSION-CENTER](http://INJECTAFER.COM/IRON-INFUSION-CENTER)**

## BEFORE YOUR FIRST INJECTAFER INFUSION

- ✓ **Call ahead** to confirm that the infusion center can provide Injectafer
- ✓ **Dress comfortably and eat as you normally would.** There are no special dietary requirements
- ✓ **Bring insurance card, photo identification,** and any other information your doctor has asked you to bring

## DURING YOUR INFUSION



Each Injectafer infusion may take about **15 minutes**



**The usual dosage is 1500 mg, administered in 2 separate doses of 750 mg each.** Your dosage may vary based on your weight

## AFTER EACH INFUSION

- ✓ Afterward, your healthcare provider will **monitor you for about 30 minutes** for signs of allergic reaction
- ✓ In some patients, an increase in blood pressure with dizziness, nausea, or flushing of the face may occur right after a dose of Injectafer. It usually goes away within 30 minutes. **Tell your doctor right away if symptoms persist or worsen**
- ✓ **Your iron stores are replenished over time.** So, it's important for you to follow up with your doctor to retest your levels and see how Injectafer is working for you

**TO ENSURE YOU RECEIVE THE COMPLETE COURSE OF TREATMENT, BE SURE TO SCHEDULE YOUR 2ND INJECTAFER TREATMENT AT LEAST 7 DAYS AFTER YOUR FIRST**

## SELECTED SAFETY INFORMATION ABOUT INJECTAFER

### What should I tell my doctor or healthcare provider before receiving Injectafer? (cont'd)

Before you receive Injectafer, tell your healthcare provider about all of your medical conditions, including if you (cont'd):

- Are pregnant or plan to become pregnant. It is not known if Injectafer will harm your unborn baby. Your healthcare provider will decide if it is safe for you to take Injectafer
- Are breastfeeding or plan to breast feed. Injectafer passes into your breast milk. It is unknown whether Injectafer would pose a risk to your baby. Talk to your healthcare provider about the best way to feed your baby during treatment with Injectafer

Tell your healthcare provider about all the medications you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**Click here to see the Full Prescribing Information and the Important Safety Information on pages 13-15.**



# injectconnect™

## SUPPORT PROGRAM

If you have an Injectafer prescription, you may be able to get help with your out-of-pocket costs.

## TWO WAYS TO HELP YOU PAY FOR INJECTAFER TREATMENTS\*

\*Restrictions apply.

# 1

### IF INJECTAFER IS COVERED BY YOUR COMMERCIAL INSURANCE, BUT YOU HAVE A CO-PAY



#### INJECTAFER SAVINGS PROGRAM

The Injectafer Savings Program may help eligible patients with their Injectafer prescription out-of-pocket responsibility.

PATIENTS RECEIVE **EACH DOSE** FOR AS LITTLE AS **\$50<sup>†</sup>**

#### For eligible patients:

- Assistance of up to \$500 per dose
- Enrollment is valid for 2 courses of treatment per 12-month period

#### HOW TO ENROLL

Ask your doctor to enroll you prior to receiving Injectafer treatment.

If your doctor cannot enroll you, you can enroll yourself in one of two ways:

- The best way to enroll is by visiting [injectafercopay.com](http://injectafercopay.com)

#### OR

- Call Daiichi Sankyo Access Central (1-866-437-4669)



# 2

### IF YOU DO NOT HAVE INSURANCE TO COVER TREATMENT OR ARE COMMERCIALY UNDERINSURED:



#### PATIENT ASSISTANCE PROGRAM

The Patient Assistance Program was created to help patients who lack health insurance or are commercially underinsured and cannot afford therapy.



Daiichi Sankyo Access Central

**1-866-4-DSI-NOW** (1-866-437-4669)

is available Monday–Friday,  
9:00 AM–8:00 PM ET

<sup>†</sup>The Injectafer Savings Program is only available for patients 1 year of age or older who are commercially insured. Insurance out of pocket must be over \$50 per dose. Additional restrictions may apply. Please see full Terms and Conditions on page 12.

**Click here to see the Full Prescribing Information and the Important Safety Information on pages 13-15.**



## Injectafer Savings Program Terms and Conditions

1. This offer is valid for commercially insured patients. Uninsured and cash-paying patients are NOT eligible for this Program.
2. Depending on insurance coverage, eligible insured patients may pay no more than \$50 per dose for two courses of treatment per 12-month period and up to a maximum savings limit of \$500 per dose, a \$1000 program limit per course of treatment. Check with your pharmacist or healthcare provider for your copay discount. Patient out-of-pocket expense may vary.
3. This offer is not valid for patients enrolled in Medicare, Medicaid, or other federal or state healthcare programs, or private indemnity or HMO insurance plans that reimburse you for the entire cost of your prescription drugs. Patients may not use this card if they are Medicare-eligible and enrolled in an employer-sponsored health plan or medical or prescription drug benefit program for retirees.
4. This offer is valid for 2 courses or 4 doses of the 750 mg dose of the Injectafer Prescription. An explanation of benefits statement must be faxed, uploaded in the portal or mailed in prior to transacting on the account numbers for co-pay assistance. One enrollment is allowed per 12-month period.
5. Daiichi Sankyo, Inc. reserves the right to rescind, revoke, or amend this offer without notice. Offer good only in the USA, including Puerto Rico, at participating pharmacies, or healthcare providers.
6. Void if prohibited by law, taxed, or restricted.
7. This account number is not transferable. The selling, purchasing, trading, or counterfeiting of this account number is prohibited by law.
8. This account number is not insurance.
9. By redeeming this account number, you acknowledge that you are an eligible patient and that you understand and agree to comply with the terms and conditions of this offer.
10. Qualified patients receiving Injectafer will be allowed a 120-day retroactive enrollment period to receive benefits under the program rules.

### Eligibility requirements for the Patient Assistance Program:

To be eligible for the program, you must

- Meet established income limits
- Lack health insurance completely or be commercially underinsured
- Be a resident of the USA or its territories, including Puerto Rico

Your provider (eg, physician, infusion center) applies to the program on your behalf.

Injectafer® (ferric carboxymaltose injection) is available by prescription only. Ask your doctor or healthcare provider if Injectafer is right for you.

### What is Injectafer?

Injectafer is a prescription iron replacement medicine administered only by or under the supervision of your healthcare provider. Injectafer is injected into your vein to treat iron deficiency anemia in adults and pediatric patients 1 year of age and older. Injectafer should be used only if you have not responded well to treatment with oral iron, or if you are intolerant to oral iron treatment. It is also used to treat iron deficiency anemia in adults with chronic kidney disease who are not receiving dialysis.

## IMPORTANT SAFETY INFORMATION

### Who should not receive Injectafer?

**You should not receive Injectafer if you** are allergic to ferric carboxymaltose or any of the other ingredients in Injectafer. The active ingredient in Injectafer is ferric carboxymaltose, the inactive ingredients are: water for injection, sodium hydroxide and/or hydrochloric acid.

### What should I tell my doctor or healthcare provider before receiving Injectafer?

Before you receive Injectafer, tell your healthcare provider about all of your medical conditions, including if you:

- Have had an allergic reaction to iron given intravenously (into your vein), including Injectafer, or to other non-oral iron treatments
- If you have, or have previously experienced, iron overload, or if your body has difficulty using iron appropriately
- Have high blood pressure
- Are pregnant or plan to become pregnant. It is not known if Injectafer will harm your unborn baby. Your healthcare provider will decide if it is safe for you to take Injectafer
- Are breastfeeding or plan to breast feed. Injectafer passes into your breast milk. It is unknown whether Injectafer would pose a risk to your baby. Talk to your healthcare provider about the best way to feed your baby during treatment with Injectafer

Tell your healthcare provider about all the medications you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**Click here to see the Full Prescribing Information and the Important Safety Information on pages 13-15.**



## How will I receive Injectafer?

Injectafer is given intravenously (into your vein) by your healthcare provider in 2 doses at least 7 days apart.

## What are the possible side effects of Injectafer?

**Injectafer can cause serious side effects, including:**

- **Serious allergic reactions** that may be life-threatening, including shock, low blood pressure, loss of consciousness, and death. Your doctor or healthcare provider will monitor you for signs and symptoms of an allergic reaction during and after each dose of Injectafer for at least 30 minutes. Other serious allergic reactions include itching, rash, hives, wheezing, or low blood pressure. You should report any signs and symptoms of an allergic reaction to Injectafer, in particular rashes, shortness of breath, and wheezing to your doctor or healthcare provider
- **High blood pressure**, sometimes with facial flushing, dizziness, or nausea, has been seen during treatment with Injectafer. This increase in blood pressure typically resolves within 30 minutes. Your doctor or healthcare provider will monitor you for signs and symptoms of an increase in blood pressure following each use of Injectafer

Other serious side effects that have been reported include rash, difficulty breathing, itching, rapid heartbeat, fever, chest discomfort, chills, swelling of the face, lips, or tongue, back pain, muscle aches, and fainting.

**The most common side effects of Injectafer include:**

- In adults: nausea, high blood pressure, flushing, pain or bruising at the injection site, skin redness, low levels of phosphorous in your blood, and dizziness. Potentially long-lasting brown staining of skin near the injection site may occur if Injectafer leaks out of the vein
- In children: low levels of phosphorous in your blood, pain or bruising at the injection site, rash, headache, and vomiting

Excessive amounts of Injectafer may lead to a condition called iron overload, which is a buildup of iron and may be harmful.

These are not all of the possible side effects of Injectafer.

Tell your doctor if you have any side effect that bothers you or that does not go away. Call your doctor for medical advice about side effects.

## General information about Injectafer

Injectafer may impact laboratory tests that measure iron in your blood for 24 hours after receiving Injectafer. Let your healthcare provider and laboratory staff know if you have received Injectafer within 24 hours of having blood tests.

**To report side effects, contact American Regent at 1-800-734-9236 or E-mail: [pv@americanregent.com](mailto:pv@americanregent.com) or Fax: 1-610-650-0170.**

**You may also report side effects to the FDA at 1-800-332-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**The risk information provided here is not comprehensive. To learn more about Injectafer, talk with your healthcare provider or pharmacist. The FDA-approved product labeling can be found at [www.injectafer.com/pdf/pi.pdf](http://www.injectafer.com/pdf/pi.pdf) or call 1-800-645-1706.**

**Please see [Full Prescribing Information](#) for Injectafer, including the bolded WARNING regarding hypersensitivity.**



**WHEN YOU'RE  
NOT SATISFIED  
WITH 10%, THERE'S  
INJECTAFER.**

**To learn more,  
visit [INJECTAFER.COM](https://www.injectafer.com)**

Injectafer® (ferric carboxymaltose injection) is available by prescription only. Ask your doctor or healthcare provider if Injectafer is right for you.

### **What is Injectafer?**

Injectafer is a prescription iron replacement medicine administered only by or under the supervision of your healthcare provider. Injectafer is injected into your vein to treat iron deficiency anemia in adults and pediatric patients 1 year of age and older. Injectafer should be used only if you have not responded well to treatment with oral iron, or if you are intolerant to oral iron treatment. It is also used to treat iron deficiency anemia in adults with chronic kidney disease who are not receiving dialysis.

**To report side effects, contact American Regent at 1-800-734-9236 or E-mail: [pv@americanregent.com](mailto:pv@americanregent.com) or Fax: 1-610-650-0170. You may also report side effects to the FDA at 1-800-332-1088 or [www.fda.gov/medwatch](https://www.fda.gov/medwatch).**

To contact us with questions or concerns about a Daiichi Sankyo product, please call us: 1-877-4DS-PROD (1-877-437-7763).

American Regent, Inc. is a member of the Daiichi Sankyo Group. Injectafer® and the Injectafer® logo are trademarks of Vifor (International) Inc., Switzerland. Injectafer® is manufactured under license from Vifor (International) Inc., Switzerland. Trademarks not owned by American Regent, Inc. or Vifor (International) are the property of their respective owners.

©2022 Daiichi Sankyo, Inc.  
03/22 PP-US-IN-2771

