

**10% OF
A BROCHURE
MAY NOT BE
ENOUGH**

**IF YOU HAVE IRON
DEFICIENCY ANEMIA...**

**10% OR LESS
OF ORAL IRON
MAY NOT BE
ENOUGH EITHER¹**



Actor Portrayal

Injectafer[®] (ferric carboxymaltose injection) is available by prescription only. Ask your doctor or healthcare provider if Injectafer is right for you.

What is Injectafer?

Injectafer is a prescription iron replacement medicine administered only by or under the supervision of your healthcare provider. Injectafer is injected into your vein to treat iron deficiency anemia in adults. Injectafer should be used only if you have not responded well to treatment with oral iron, or if you are intolerant to oral iron treatment. It is also used to treat iron deficiency anemia in adults with chronic kidney disease who are not receiving dialysis.

It is not known if Injectafer is safe and effective for use in children.

SELECTED SAFETY INFORMATION

Who should not receive Injectafer?

You should not receive Injectafer if you are allergic to ferric carboxymaltose or any of the other ingredients in Injectafer. The active ingredient in Injectafer is ferric carboxymaltose, the inactive ingredients are: water for injection, sodium hydroxide and/or hydrochloric acid.

[Click here](#) to see the Full Prescribing Information, and see Important Safety Information on pages 20-21.



WELCOME TO ANOTHER WAY TO TREAT IRON DEFICIENCY ANEMIA (IDA)²



SELECTED SAFETY INFORMATION ABOUT INJECTAFER

What is Injectafer?

Injectafer is a prescription iron replacement medicine administered only by or under the supervision of your healthcare provider. Injectafer is injected into your vein to treat iron deficiency anemia in adults. Injectafer should be used only if you have not responded well to treatment with oral iron, or if you are intolerant to oral iron treatment. It is also used to treat iron deficiency anemia in adults with chronic kidney disease who are not receiving dialysis.

It is not known if Injectafer is safe and effective for use in children.

Considering Injectafer

This brochure will give you information about IDA. In this section, you'll learn about the challenges of oral iron supplements, and about another treatment option you may want to discuss with your doctor.^{1,2}

Getting Started on Injectafer

If your doctor has prescribed IV iron, the information in this section will help you get started on Injectafer, the only IV iron that can provide up to 1500 mg of iron in just 2 administrations of 750 mg. You will learn how and where Injectafer is administered and what to expect.²

Savings & Support


Only Injectafer offers the InjectConnect Support Program. Find out if you're eligible and learn how to enroll.

[Click here to see the Full Prescribing Information, and see Important Safety Information on pages 20-21.](#)

WHAT IS IDA?

- **Iron deficiency anemia (IDA)** is a condition that interferes with the formation and function of red blood cells³
- **Red blood cells contain hemoglobin**, which is a protein that helps carry oxygen from the lungs to cells in the body³
- When you don't have enough healthy red blood cells to carry enough oxygen to cells throughout the body, **that's anemia**⁴
- **IDA is the most common form of anemia**³

WHAT ARE SOME OF THE SYMPTOMS OF IDA?^{5*}




*Some patients with IDA may not experience symptoms. As lab markers are the only way to diagnose IDA, it is important that patients follow up with their doctor for lab work.²

Injectafer has not been studied or approved for treating symptoms of IDA.

WHAT CAUSES IDA?

- **Low iron levels**—an iron deficiency—cause IDA³
- IDA occurs when iron in your body **becomes depleted or you aren't adequately replenishing** what your body uses³
- Your iron deficiency is the **difference between the amount of iron you have and the amount you need** to keep you in good health^{3,6}
- **If your iron deficiency progresses, you may have IDA**⁶

IDA CAN COEXIST WITH MANY DIFFERENT DISEASES OR CONDITIONS



YOU MAY HAVE IRON DEFICIENCY ANEMIA IF YOU DO NOT HAVE ENOUGH IRON IN YOUR BODY³
INJECTAFER MAY BE AN OPTION TO REPLACE THE IRON YOUR BODY IS MISSING²

SELECTED SAFETY INFORMATION ABOUT INJECTAFER

Who should not receive Injectafer?

You should not receive Injectafer if you are allergic to ferric carboxymaltose or any of the other ingredients in Injectafer. The active ingredient in Injectafer is ferric carboxymaltose, the inactive ingredients are: water for injection, sodium hydroxide and/or hydrochloric acid.

[Click here to see the Full Prescribing Information, and see Important Safety Information on pages 20-21.](#)



WHY CAN'T I JUST TAKE AN IRON PILL?

THERE ARE SEVERAL REASONS WHY PILLS MAY NOT BE SUITABLE FOR SOME PEOPLE WITH IDA.¹

- Oral iron supplements can cause hard-to-tolerate **side effects**¹
- **Poor absorption**—the digestive tract is only able to absorb a small portion of the iron in an oral iron supplement, so your body may not get the full dose of iron needed from pills¹



IF ORAL IRON IS INADEQUATE OR NOT WELL TOLERATED, IV IRON MAY BE AN OPTION FOR YOU.²

- With IV treatments, **iron is delivered directly into the bloodstream** through a vein^{2,9}
- IV iron isn't processed by the digestive system, so **100% is delivered into the bloodstream**²
- Getting IV iron is a process known as infusion⁹

The information provided by Daiichi Sankyo, Inc. is not intended to replace your healthcare provider's medical advice.



COMMON LAB MARKERS FOR MONITORING IDA AND NORMAL VALUES IN ADULTS*

Your iron deficiency is determined by lab markers in your blood. If your lab values are below the normal range, it's important to talk with your doctor about the best option for restoring your iron levels.¹

	Male	Female
Hemoglobin (Hb)	13.5 - 17.5 g/dL	12.0 - 15.5 g/dL ¹⁰
Ferritin	40 - 300 µg/L	20 - 200 µg/L ¹¹
Transferrin saturation (TSAT)	20% to 50%	20% to 50% ¹²

An additional test for total iron binding capacity (TIBC) values may be required to diagnose IDA.⁵ Normal values for TIBC range from 240 to 450 µg/dL in healthy patients.¹³ Please note that TIBC values were not measured in Injectafer clinical trials.

*Normal values can vary for many reasons, including conditions you may have or where your lab work was done. That's why it's important to discuss your lab results with your doctor to find out what they mean for you.

SPEAK WITH YOUR DOCTOR OR HEALTHCARE PROFESSIONAL IF YOU WOULD LIKE TO LEARN MORE ABOUT INJECTAFER.

SELECTED SAFETY INFORMATION ABOUT INJECTAFER

What should I tell my doctor or healthcare provider before receiving Injectafer?

Before you receive Injectafer, tell your healthcare provider about all of your medical conditions, including if you:

- Have had an allergic reaction to iron given intravenously (into your vein), including Injectafer, or to other non-oral iron treatments
- If you have, or have previously experienced, iron overload, or if your body has difficulty using iron appropriately
- Have high blood pressure
- Are pregnant or plan to become pregnant. It is not known if Injectafer will harm your unborn baby. Your healthcare provider will decide if it is safe for you to take Injectafer
- Are breastfeeding or plan to breast feed. Injectafer passes into your breast milk. It is unknown whether Injectafer would pose a risk to your baby. Talk to your healthcare provider about the best way to feed your baby during treatment with Injectafer

[Click here to see the Full Prescribing Information, and see Important Safety Information on pages 20-21.](#)




HOW CAN INJECTAFER HELP?



Injectafer is indicated for use in patients who have intolerance to or have had an unsatisfactory response to oral iron.²

INJECTAFER IS THE ONLY IV IRON THAT PROVIDES UP TO 1500 MG OF IRON IN JUST 2 ADMINISTRATIONS—UP TO 750 MG EACH, SEPARATED BY AT LEAST 7 DAYS²

- Injectafer is for adults who **can't tolerate oral iron or who don't respond well to it**²
- Unlike with pills, **100% of the iron** in Injectafer goes directly into the bloodstream²
- Proven in clinical studies in more than 8800 patients, **Injectafer demonstrated greater increases in hemoglobin** at follow-up than either oral iron or another IV iron^{2,15}
- Injectafer treatments are typically administered at an infusion center, and **treatments may take only 15 minutes**, followed by a period of around 30 minutes, during which you will be monitored for possible side effects^{2,9}
- **Potential side effects** with Injectafer include, but are not limited to, allergic reactions, nausea, temporary hypertension, and flushing² (see list of side effects on page 21)



More than 800,000 people have been treated with Injectafer in the United States alone.¹⁶

Not an actual patient.

Based on IQVIA data through December 2018.

SELECTED SAFETY INFORMATION ABOUT INJECTAFER

- Are breastfeeding or plan to breast feed. It is unknown whether Injectafer would pose a risk to your baby. Talk to your healthcare provider about the best way to feed your baby during treatment with Injectafer.

What are the possible side effects of Injectafer?

Injectafer can cause serious side effects, including:

- **Serious allergic reactions** that may be life-threatening, including shock, low blood pressure, loss of consciousness, and death. Your doctor or healthcare provider will monitor you for signs and symptoms of an allergic reaction during and after each dose of Injectafer for at least 30 minutes. Other serious allergic reactions include itching, rash, hives, wheezing, or low blood pressure. You should report any signs and symptoms of an allergic reaction to Injectafer, in particular rashes, shortness of breath and wheezing to your doctor or healthcare provider.
- **High blood pressure**, sometimes with facial flushing, dizziness, or nausea, has been seen during treatment with Injectafer. This increase in blood pressure typically resolves within 30 minutes. Your doctor or healthcare provider will monitor you for signs and symptoms of an increase in blood pressure following each use of Injectafer.

Other serious side effects that have been reported include rash, difficulty breathing, itching, rapid heartbeat, fever, chest discomfort, chills, swelling of the face, lips, or tongue, back pain, muscle aches, and fainting.

[Click here to see the Full Prescribing Information, and see Important Safety Information on pages 20-21.](#)



READY TO GET STARTED ON INJECTAFER?

Your doctor has prescribed Injectafer for you because it may be an effective way to replace the iron you need. Injectafer is the only IV iron that provides up to 1500 mg of iron in just 2 doses. Each dose provides up to 750 mg of iron, and doses are separated by at least 7 days.²

WHERE IS INJECTAFER GIVEN?



- **Injectafer is given by IV infusion**, usually at an infusion center, where a healthcare professional can administer it directly into the bloodstream through a vein⁹

- Infusion centers are medical facilities equipped and staffed for **administering infusions**⁹



- Each Injectafer infusion may take about **15 minutes**²

- Afterward, your healthcare provider will monitor you for about 30 minutes for signs of allergic reaction²



Find an infusion center that provides Injectafer by using the Injectafer Infusion Center Locator at www.injectafer.com/iron-infusion-center

BEFORE YOUR FIRST INJECTAFER INFUSION

- ✓ **Call ahead** to confirm that the infusion center can provide Injectafer IV iron
- ✓ **Ask about Injectafer by name.** Injectafer is the only IV iron that can provide up to 1500 mg of iron in just 2 doses of 750 mg²
- ✓ **Dress comfortably and eat as you normally would.** There are no special dietary requirements
- ✓ **Bring insurance coverage identification** and any other information your doctor has asked you to bring
- ✓ Injectafer offers co-pay support through the **InjectConnect Support Program*** (see page 16 to learn more)

*For eligible patients. Restrictions apply.

SELECTED SAFETY INFORMATION ABOUT INJECTAFER

What are the possible side effects of Injectafer? (continued)

The most common side effects of Injectafer include:

- Nausea, high blood pressure, flushing, low levels of phosphorus in your blood, dizziness, vomiting, headache, an increase in certain liver enzymes, and pain or bruising at the injection site. Potentially long-lasting brown staining of skin near the injection site may occur if Injectafer leaks out of the vein.

Excessive amounts of Injectafer may lead to a condition called iron overload, which is a buildup of iron and may be harmful.

These are not all of the possible side effects of Injectafer.

Tell your doctor if you have any side effect that bothers you or that does not go away. Call your doctor for medical advice about side effects.

[Click here to see the Full Prescribing Information, and see Important Safety Information on pages 20-21.](#)



GETTING YOUR INJECTAFER INFUSIONS

DURING YOUR INFUSION

- **The usual dosage is 1500 mg, administered in 2 separate doses of 750 mg each.** Your dosage may vary based on your weight²
- **Your infusion may take about 15 minutes,** and your doctor or nurse will monitor your response during and after your infusion²

AFTER EACH INFUSION

- **Your iron stores are replenished over time. So it's important for you to follow up with your doctor and retest your levels¹⁷**
- In some patients, an increase in blood pressure with dizziness, nausea, or flushing of the face may occur right after a dose of Injectafer. It usually goes away within 30 minutes. **Tell your doctor right away if symptoms persist or worsen²**
- **Be sure to follow up with your physician** to see how Injectafer is working for you
- Track your treatments and progress **using the progress tracker in this brochure**

SELECTED SAFETY INFORMATION ABOUT INJECTAFER

General information about Injectafer

Injectafer may impact laboratory tests that measure iron in your blood for 24 hours after receiving Injectafer. Let your healthcare provider and laboratory staff know if you have received Injectafer within 24 hours of having blood tests.

BE SURE TO COMPLETE YOUR INJECTAFER COURSE OF TREATMENT

REMEMBER, IT TAKES TWO DOSES

- **Be sure to come back for your second dose** after at least 7 days²
- Remember to **schedule or confirm your next appointment**
- **Track your treatments and progress** using the progress tracker in this brochure

$$750 \text{ mg} + 750 \text{ mg} = 1500 \text{ mg}^2$$

WHAT SIDE EFFECTS COULD OCCUR IN THE FOLLOWING DAYS?

- Some patients experience nausea, high blood pressure, flushing, low levels of phosphorous in their blood, dizziness, vomiting, headache, an increase in certain liver enzymes, and pain or bruising at the injection site during or immediately after an infusion²
- These symptoms may persist or show up in the following hours or days²
- These are not all of the possible side effects of Injectafer. Call your doctor for medical advice about side effects²

SEEK MEDICAL ATTENTION IF MORE SERIOUS SIDE EFFECTS ARISE²

Serious side effects may include, but are not limited to:

Allergic reactions including itching, hives, wheezing, low blood pressure, and high blood pressure, sometimes with face flushing, dizziness, or nausea.

[Click here to see the Full Prescribing Information, and see Important Safety Information on pages 20-21.](#)



INJECTAFER TREATMENT REMINDERS



	Dose #1	Dose #2
Date		
Time		
Infusion Center Location		

Doctor follow-up:

Date	Time
Location	

Baseline iron levels:

Date	Hemoglobin	TSAT	Ferritin
------	------------	------	----------

Iron levels at follow-up:

Date	Hemoglobin	TSAT	Ferritin
------	------------	------	----------

GET THE LATEST INFORMATION ABOUT IDA AND INJECTAFER.

- To learn more about IDA and Injectafer, visit www.injectafer.com/info
- For news about Injectafer, sign up for the Infuse News email updates: visit www.injectafer.com/signup

WHAT IS INJECTAFER?

Injectafer is a prescription iron replacement medicine administered only by or under the supervision of your healthcare provider. Injectafer is injected into your vein to treat iron deficiency anemia in adults. Injectafer should be used only if you have not responded well to treatment with oral iron, or if you are intolerant to oral iron treatment. It is also used to treat iron deficiency anemia in adults with chronic kidney disease who are not receiving dialysis.

It is not known if Injectafer is safe and effective for use in children.

[Click here to see the Full Prescribing Information, and see Important Safety Information on pages 20-21.](#)



DIET AND LIFESTYLE TIPS TO HELP KEEP YOUR IRON LEVELS HEALTHY

DISCUSS A DIET PLAN WITH YOUR DOCTOR WHO MAY SUGGEST THAT YOU EAT ONE OR MORE OF THESE IRON-RICH FOODS⁵

- Red meat: beef, pork, lamb
- Liver
- Poultry, especially dark meat
- Dark leafy greens: broccoli, kale, turnip greens, collard greens
- Legumes: beans and peas
- Iron-enriched grains, pasta, rice, cereals



REDUCE ALCOHOL CONSUMPTION

Alcohol may reduce the body's ability to make red blood cells and may shorten red blood cell lifespan.¹⁸

KEEP ACTIVE

Remember to speak to your healthcare provider before you begin any fitness program.

RELAX

Don't be afraid to ask for help, understanding, and patience when needed.

The tips included in this brochure are meant to aid, not replace, conversations with your doctor. Please talk to a healthcare professional to determine the best health tips for you.

injectconnect™

SUPPORT PROGRAM

TWO WAYS TO HELP YOU PAY FOR INJECTAFER TREATMENTS*

*Restrictions apply.

1

If Injectafer is covered by your commercial insurance, but you have a co-pay:



INJECTAFER SAVINGS PROGRAM

Injectafer is the only IV iron with a copay assistance program.†

If you're eligible,‡ you may:

- Receive assistance of up to \$500 per dose
- Receive a maximum benefit of up to \$1000 per course of treatment (2 doses)
- Re-enroll once during a 12-month period if you need another course of treatment

FIRST DOSE
for as little as

\$50

Up to 750 mg of iron for qualified patients†

SECOND DOSE
for as little as

\$0

Up to 750 mg of iron for qualified patients†

HOW TO ENROLL



To find out if you're eligible, or to enroll, apply online at www.injectafer.com. You will be asked to provide information about yourself, your doctor, and your current healthcare plan. You can also call the IV Iron Hotline to learn more.

2

If you do not have insurance to cover treatment:



PATIENT ASSISTANCE PROGRAM

The Patient Assistance Program was created to help patients who lack health insurance and cannot afford therapy.

Call the IV Iron Hotline to find out if you are eligible.

(See page 19 for eligibility requirements.)



Our toll-free IV Iron Hotline
1-877-4-IV-IRON (1-877-448-4766)
is available Monday–Friday, 9:00 AM–8:00 PM ET

†As of November 2019.

‡The Injectafer Savings Program is only available for adults 18 years or older who are commercially insured or cash-paying patients. The program provides up to a maximum savings limit of \$500 per dose and a maximum benefit of up to \$1000 per course of treatment (2 doses). Enrollment is valid for 2 courses of therapy (4 doses). Insurance out of pocket must be over \$50. Additional restrictions may apply. Please see full Terms and Conditions on page 16.

Click here to see the Full Prescribing Information, and see Important Safety Information on pages 20-21.

Terms and Conditions for the Injectafer Savings Program:

1. This offer is valid for commercially-insured as well as cash paying patients.
2. Depending on insurance coverage, eligible insured patients may pay no more than \$50 for Injectafer for the first dose and \$0 for Injectafer for the second dose, up to a maximum savings limit of \$500 per dose, a \$1,000 program limit per course of therapy. Check with your pharmacist or healthcare provider for your copay discount. Patient out-of-pocket expense may vary.
3. This offer is not valid for patients enrolled in Medicare, Medicaid, or other federal or state healthcare programs, or private indemnity or HMO insurance plans that reimburse you for the entire cost of your prescription drugs. Patients may not use this card if they are Medicare-eligible and enrolled in an employer-sponsored health plan or medical or prescription drug benefit program for retirees.
4. The offer is valid for 2 courses, or 4 doses, of an Injectafer prescription. An explanation of benefits statement must be faxed in prior to transacting on the account numbers for assistance. One enrollment is allowed per 12-month period.
5. Daiichi Sankyo, Inc. reserves the right to rescind, revoke, or amend this offer without notice.
6. Offer good only in the USA, including Puerto Rico, at participating pharmacies or healthcare providers.
7. Void if prohibited by law, taxed, or restricted.
8. This account number is not transferable. The selling, purchasing, trading, or counterfeiting of this account number is prohibited by law.
9. This account number is not insurance.
10. By redeeming this account number, you acknowledge that you are an eligible patient and that you understand and agree to comply with the terms and conditions of this offer.
11. Qualified patients receiving Injectafer will be allowed a 60-day retroactive enrollment period to receive benefits under the program rules.

Eligibility Requirements for the Patient Assistance Program:

To be eligible for the program, you must:

- Completely lack health insurance and be ineligible for public insurance or financing
- Be a US citizen, legal entrant in the United States, or permanent resident. Proof of citizenship or legal residency may also be required
- Meet income and other criteria

Providers (hospitals, physicians, or infusion centers) must apply to the program on behalf of their patients. They will need to fill out and fax the Patient Assistance Request form and Enrollment Application.

REFERENCES

1. Zhu A, Kaneshiro M, Kaunitz JD. Evaluation and treatment of iron deficiency anemia: a gastroenterological perspective. *Dig Dis Sci*. 2010;55(3):548-559.
2. Injectafer® [package insert]. Shirley, NY: American Regent, Inc.; 2018.
3. Anemia. American Society of Hematology website. <http://www.hematology.org/Patients/Anemia>. Accessed July 26, 2018.
4. Anemia. Mayo Clinic website. <https://www.mayoclinic.org/diseases-conditions/anemia/symptoms-causes/syc-20351360>. Accessed August 31, 2018.
5. Iron-Deficiency Anemia. American Society of Hematology website. <http://www.hematology.org/Patients/Anemia/Iron-Deficiency>. Accessed August 14, 2018.
6. Bermejo F, García-López S. A guide to diagnosis of iron deficiency and iron deficiency anemia in digestive disease. *World J Gastroenterol*. 2009;15(37):4638-4643.
7. Anemia in chronic kidney disease. National Institute of Diabetes and Digestive and Kidney Diseases website. <https://www.niddk.nih.gov/health-information/kidney-disease/anemia>. Accessed August 31, 2018.
8. Okonko DO, Mandal AK, Missouri CG, Poole-Wilson PA. Disordered iron homeostasis in chronic heart failure: prevalence, predictors, and relation to anemia, exercise capacity, and survival. *J Am Coll Cardiol*. 2011;58(12):1241-1251.
9. National Infusion Center Association. I Infusion therapy: frequently asked questions. <https://infusioncenter.org/faqs>. Accessed August 14, 2018.
10. Hemoglobin test. Mayo Clinic website. <https://www.mayoclinic.org/tests-procedures/hemoglobin-test/about/pac-20385075>. Accessed August 6, 2018.
11. Iron-deficiency anemia. National Heart, Lung, and Blood Institute website. <https://www.nhlbi.nih.gov/health-topics/iron-deficiency-anemia>. Accessed August 6, 2018.
12. Transferrin saturation. Medscape website. <http://emedicine.medscape.com/article/2087960-overview>. Accessed August 31, 2018.
13. Total iron binding capacity. MedlinePlus website. <https://medlineplus.gov/ency/article/003489.htm>. Accessed June 26, 2019.
14. Onken JE, Bregman DB, Harrington RA, et al. A multicenter, randomized, active-controlled study to investigate the efficacy and safety of intravenous ferric carboxymaltose in patients with iron deficiency anemia. *Transfusion*. 2014;54(2):306-315.
15. Koch TA, Myers J, Goodnough LT. Intravenous iron therapy in patients with iron deficiency anemia: dosing considerations. *Anemia*. 2015;763576. doi:10.1155/2015/763576.
16. Data on file. Daiichi Sankyo Inc., Basking Ridge, NJ.
17. Nall R. Iron infusion: uses, benefits, and what to expect. Medical News Today website. <https://www.medicalnewstoday.com/articles/318827.php>. Accessed August 31, 2018.
18. Ballard HS. The hematological complications of alcoholism. *Alcohol Health Res World*. 1997;21(1):42-52.

Click here to see the Full Prescribing Information, and see Important Safety Information on pages 20-21.



ABOUT INJECTAFER

Injectafer® (ferric carboxymaltose injection) is available by prescription only. Ask your doctor or healthcare provider if Injectafer is right for you.

What is Injectafer?

Injectafer is a prescription iron replacement medicine administered only by or under the supervision of your healthcare provider. Injectafer is injected into your vein to treat iron deficiency anemia in adults. Injectafer should be used only if you have not responded well to treatment with oral iron, or if you are intolerant to oral iron treatment. It is also used to treat iron deficiency anemia in adults with chronic kidney disease who are not receiving dialysis.

It is not known if Injectafer is safe and effective for use in children.

IMPORTANT SAFETY INFORMATION

Who should not receive Injectafer?

You should not receive Injectafer if you are allergic to ferric carboxymaltose or any of the other ingredients in Injectafer. The active ingredient in Injectafer is ferric carboxymaltose, the inactive ingredients are: water for injection, sodium hydroxide and/or hydrochloric acid.

What should I tell my doctor or healthcare provider before receiving Injectafer?

Before you receive Injectafer, tell your healthcare provider about all of your medical conditions, including if you:

- Have had an allergic reaction to iron given intravenously (into your vein), including Injectafer, or to other non-oral iron treatments
- If you have, or have previously experienced, iron overload, or if your body has difficulty using iron appropriately
- Have high blood pressure
- Are pregnant or plan to become pregnant. It is not known if Injectafer will harm your unborn baby. Your healthcare provider will decide if it is safe for you to take Injectafer
- Are breastfeeding or plan to breast feed. Injectafer passes into your breast milk. It is unknown whether Injectafer would pose a risk to your baby. Talk to your healthcare provider about the best way to feed your baby during treatment with Injectafer.

Tell your healthcare provider about all the medications you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of Injectafer?

Injectafer can cause serious side effects, including:

- **Serious allergic reactions** that may be life-threatening, including

shock, low blood pressure, loss of consciousness, and death. Your doctor or healthcare provider will monitor you for signs and symptoms of an allergic reaction during and after each dose of Injectafer for at least 30 minutes. Other serious allergic reactions include itching, rash, hives, wheezing, or low blood pressure. You should report any signs and symptoms of an allergic reaction to Injectafer, in particular rashes, shortness of breath and wheezing to your doctor or healthcare provider.

- **High blood pressure**, sometimes with facial flushing, dizziness, or nausea, has been seen during treatment with Injectafer. This increase in blood pressure typically resolves within 30 minutes. Your doctor or healthcare provider will monitor you for signs and symptoms of an increase in blood pressure following each use of Injectafer.

Other serious side effects that have been reported include rash, difficulty breathing, itching, rapid heartbeat, fever, chest discomfort, chills, swelling of the face, lips, or tongue, back pain, muscle aches, and fainting.

The most common side effects of Injectafer include:

- Nausea, high blood pressure, flushing, low levels of phosphorous in your blood, dizziness, vomiting, headache, an increase in certain liver enzymes, and pain or bruising at the injection site. Potentially long-lasting brown staining of skin near the injection site may occur if Injectafer leaks out of the vein.

Excessive amounts of Injectafer may lead to a condition called iron overload, which is a buildup of iron and may be harmful.

These are not all of the possible side effects of Injectafer.

Tell your doctor if you have any side effect that bothers you or that does not go away. Call your doctor for medical advice about side effects.

General information about Injectafer

Injectafer may impact laboratory tests that measure iron in your blood for 24 hours after receiving Injectafer. Let your healthcare provider and laboratory staff know if you have received Injectafer within 24 hours of having blood tests.

To report side effects, contact American Regent at 1-800-734-9236 or E-mail: pv@americanregent.com or Fax: 1-610-650-0170.

You may also report side effects to the FDA at 1-800-332-1088 or www.fda.gov/medwatch.

The risk information provided here is not comprehensive. To learn more about Injectafer, talk with your healthcare provider or pharmacist. The FDA-approved product labeling can be found at www.injectafer.com/pdf/pi.pdf or call 1-800-645-1706.

Please see Full Prescribing Information for Injectafer.

[Click here to see the Full Prescribing Information.](#)



IF YOU HAVE IRON DEFICIENCY ANEMIA AND ORAL IRON SUPPLEMENTS AREN'T WORKING WELL ENOUGH...

- ✓ Ask your doctor about IV iron with Injectafer²
- ✓ Find out if you're eligible for **cost savings with InjectConnect***
- ✓ **Call ahead** and make sure the infusion center can provide Injectafer
- ✓ Be sure to complete your full course of treatment—
2 doses of 750 mg = 1500 mg^{2†}

*Restrictions apply.

[†]For patients weighing less than 50 kg (110 lb): Give Injectafer in two doses separated by at least 7 days and give each dose as 15 mg/kg body weight.

Injectafer[®] (ferric carboxymaltose injection) is available by prescription only. Ask your doctor or healthcare provider if Injectafer is right for you.

What is Injectafer?

Injectafer is a prescription iron replacement medicine administered only by or under the supervision of your healthcare provider. Injectafer is injected into your vein to treat iron deficiency anemia in adults. Injectafer should be used only if you have not responded well to treatment with oral iron, or if you are intolerant to oral iron treatment. It is also used to treat iron deficiency anemia in adults with chronic kidney disease who are not receiving dialysis.

It is not known if Injectafer is safe and effective for use in children.

To report side effects, contact American Regent at 1-800-734-9236 or E-mail: pv@americanregent.com or Fax: 1-610-650-0170. You may also report side effects to the FDA at 1-800-332-1088 or www.fda.gov/medwatch.

To contact us with questions or concerns about a Daiichi Sankyo product, please call us: 1-877-4DS-PROD (1-877-437-7763).

American Regent, Inc. is a member of the Daiichi Sankyo Group. Injectafer[®] and the Injectafer[®] logo are trademarks of Vifor (International) Inc., Switzerland. Injectafer[®] is manufactured under license from Vifor (International) Inc., Switzerland. Trademarks not owned by American Regent, Inc. or Vifor (International) are the property of their respective owners.

©2019 Daiichi Sankyo, Inc. PP-US-IN-1215 11/19

[Click here](#) to see the Full Prescribing Information, and see Important Safety Information on pages 20-21.

